


What Climate Change Means for Montana

Montana's climate is changing. In the past century, most of the state has warmed about two degrees (F). Heat waves are becoming more common, and snow is melting earlier in spring. Rising temperatures and recent droughts have killed many trees by drying out soils, increasing the risk of forest fires, or enabling outbreaks of forest insects. In the coming decades, the changing climate is likely to decrease the availability of water in Montana, affect agricultural yields, and further increase the risk of wildfires.

The climate is changing because Earth is warming. People have increased the amount of carbon dioxide in the air by 40 percent since the late 1700s. Other heat-trapping greenhouse gases are also increasing. These gases have warmed the surface and lower atmosphere of our planet about one degree during the last 50 years. Evaporation increases as the atmosphere warms, which increases humidity, average rainfall, and the frequency of heavy rainstorms in many places—but contributes to drought in others.


Greenhouse gases are also changing the world's oceans and ice cover. Carbon dioxide reacts with water to form carbonic acid, so the oceans are becoming more acidic. The surface of the ocean has warmed about one degree during the last 80 years, and sea level is rising at an increasing rate. Warming is causing snow to melt earlier in spring.



Rising temperatures in the last century. Montana has warmed more than most of the contiguous United States. Source: EPA, *Climate Change Indicators in the United States*.

Snowpack and Glaciers

As the climate warms, less precipitation falls as snow, and more snow melts during winter. That decreases snowpack—the amount of snow that accumulates over the winter. Since the 1950s, the snowpack in Montana has been decreasing. Diminishing snowpack can shorten the season for skiing and other forms of winter tourism and recreation. The tree line may shift, as higher temperatures and a longer season without snow on the ground allow subalpine fir and other high-altitude trees to grow at higher elevations. A higher tree line would decrease the extent of alpine tundra ecosystems, which could threaten some species.



Trends in April snowpack in Montana, 1955–2013. The snowpack has declined at most monitoring sites in Montana. Source: EPA.

More than one thousand glaciers cover about 26 square miles of mountains in Montana, but that area is decreasing in response to rising temperatures. Glacier National Park's glaciers receded rapidly during the last century. Several of these glaciers are likely to disappear by 2030 if current trends continue. Areas that are no longer covered by glaciers may still accumulate snowpack, but the snow will no longer remain year-round.



Repeat photographs of Sperry Glacier in Glacier National Park. Source: USGS.

Precipitation and Water Resources

Changing the climate is likely to increase the demand for water and make it more available. Warmer temperatures increase evaporation and water use by plants. Increases in rainfall, however, are likely to offset these losses so that soil moisture increases slightly or remains about the same as today. More water is likely to run off into the upper Missouri River and its tributaries.

In areas that depend on melting snow, however, the supply of water is likely to decline. Mountain snowpacks are natural reservoirs that collect the snow that falls during winter and release water when the snow melts during spring and summer. Dams capture meltwater and retain it for use later in the year. But upstream of these dams, as the snowpack declines, less water is available during droughts for ecosystems, water-based recreation, and landowners who draw water directly from a natural lake or flowing river.

Agriculture

Rising temperatures and changes in rainfall are likely to have both positive and negative effects on Montana's farms and ranches, and the net effect is unknown. Higher temperatures reduce yields of wheat, but higher concentrations of carbon dioxide are likely to increase yields by a similar amount. Warmer and shorter winters may allow for a longer growing season, which could allow two crops per year instead of one in some instances. But warmer winters may also promote the growth of weeds and pests.

Rising carbon dioxide concentrations are likely to increase the productivity of rangelands. Provided that the quality of forage does not deteriorate, the higher range productivity would increase cattle production.

Warmer winters could also benefit ranches by reducing losses to winter storms. During the winter of 1996–1997, for example, high winds and heavy snow killed half of the newborn calves and 100,000 adult cows in the Northern Great Plains. But warmer summers would at least partly offset the benefit of warmer winters, because hot weather causes cows to eat less and grow more slowly, and it can threaten their health. Over the next 70 years, the number of days above 100°F in Montana is likely to double.

Wildfires

Higher temperatures and drought are likely to increase the severity, frequency, and extent of wildfires in Montana, which could harm property, livelihoods, and human health. On average, about 2 percent of the land in the state has burned per decade since 1984. Wildfire smoke pollutes the air and can increase medical visits for respiratory and heart problems.



Firefighters battle the Taylor Creek blaze in southeastern Montana in 2012. Credit: Gerald Vickers, National Wildfire Coordinating Group.

Forests

Longer growing seasons and increased carbon dioxide concentrations could increase the productivity of forests, but warmer conditions also make forests more susceptible to pests. Temperature controls the life cycle and winter mortality rates of pests such as bark beetles, which have infested millions of acres and killed millions of trees across the West in recent decades. With higher winter temperatures, some pests can persist year-round, and new pests and diseases may become established. Drought also reduces the ability of trees to mount a defense against attacks from beetles and other pests.

Human Health

Extremely hot and cold days can be unhealthy—even dangerous. Certain people are especially vulnerable, including children, the elderly, the sick, and the poor. The elderly may be particularly prone to heat stress and other heat-related health problems, including dehydration, cardiovascular strain, and respiratory problems. Those with low incomes may be particularly vulnerable if they lack air conditioning. Power failures due to severe weather can also present risks, especially in lightly populated areas where access to the necessary support services may be limited. While these risks will increase as the climate becomes warmer, illnesses and deaths due to cold weather and snow are likely to decline.